



WRF sensitivity experiments for the mesoscale NEWA wind atlas production run

Hahmann, Andrea N.; Witha, Björn; Sile, Tija; Dorenkaemper, Martin; Söderberg, Stefan; Navarro, Jorge ; Leroy, Grégoire; Folch, Arnau; Bustamante, Elena Garcia ; Gonzalez-Rouco, Fidel

Publication date:
2018

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Hahmann, A. N. (Author), Witha, B. (Author), Sile, T. (Author), Dorenkaemper, M. (Author), Söderberg, S. (Author), Navarro, J. (Author), Leroy, G. (Author), Folch, A. (Author), Bustamante, E. G. (Author), & Gonzalez-Rouco, F. (Author). (2018). WRF sensitivity experiments for the mesoscale NEWA wind atlas production run. Sound/Visual production (digital)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

WRF sensitivity experiments for the mesoscale NEWA wind atlas production run

Andrea N. Hahmann (ahah@dtu.dk)¹, Björn Witha², Tija Sile³, Martin Dörenkaemper⁴, Stefan Söderberg⁵, Jorge Navarro⁶, Grégoire Leroy⁷, Arnau Folch⁸, Elena Garcia Bustamante⁹, and Fidel Gonzalez-Rouco⁹


¹DTU, Wind Energy Department, Denmark (ahah@dtu.dk), ²ForWind, Oldenburg University, Germany, ³University of Latvia, Latvia, ⁴Fraunhofer IWES, Germany, ⁵WeatherTech, Sweden, ⁶CIEMAT, Spain, ⁷3E, Belgium, ⁸BSC, Spain, ⁹UCM, Spain



The New European Wind Atlas (NEWA) project

Objectives:



- **Accurate mapping of wind conditions for the estimation of wind resources** and loads
- **Development and testing of the “model-chain”**
- A series of field atmospheric experiments to validate the models and the final atlas




<http://www.neweuropeanwindatlas.eu>

Green – NEWA member countries


Red – field experiments sites



Find the best set of model parameterizations




First test of choices



Find the optimal domain size and arrangement

Mean wind speed at 100 m AGL



Phase0: Sensitivity experiments


Phase0: NEWA WRF sensitivity experiments

Objectives:

- Homogenize the expertise of the various participating groups
- Investigate model sensitivity and whether it is homogeneous across different European wind climates
- Year-long (2015) simulations for five different regions in Europe
- Two PBL parameterizations and
- Two integration methods

PBL	Method
MYNN	36 hours simulation, 12 hours spin-up/
YSU	8 days simulation, 1 day spin-up, nudging D1

MYNN - daily	MYNN - weekly
YSU - daily	YSU - weekly



Annual mean wind speed
(m/s) difference:


$$\bar{U}_{\text{MYNN}} - \bar{U}_{\text{YSU}}$$

Daily runs

h=100m



Mean simulated wind

lev: 100 m




Mean simulated wind

lev: 100 m




Mean simulated wind

lev: 100 m





Mean simulated wind

lev: 100 m



Mean simulated wind

lev: 100 m











Annual
mean wind
speed (m/s)
difference:

$$\bar{U}_{\text{MYNN}} - \bar{U}_{\text{YSU}}$$

Daily runs

h=100m

Surface
roughness
length (m)




Annual
mean wind
speed (m/s)
difference:



$$\bar{U}_{\text{MYNN}} - \bar{U}_{\text{YSU}}$$

Daily runs

$h=100\text{m}$




Surface
roughness
length (m)



Unstable fraction (%)

Unstable ($1/L < -0.005$) %




Annual mean wind speed (m/s) difference:

$$\bar{U}_{\text{MYNN}} - \bar{U}_{\text{YSU}}$$

Daily runs

h=100m



From Monin-Obukhov similarity theory:



$$U(z) = \frac{u_*}{\kappa} [\ln(z/z_0) - \psi(z/L)]$$

Wind shear is a function of surface roughness and atmospheric stability

Thus, the "first-order" response of the mean wind speed to the PBL scheme depends on z_0 and $1/L$




Surface roughness length (m)



DTU Wind Energy, Technical University of Denmark

Unstable fraction (%)



4/26/2018

Phase1: Model evaluation

Phase1: Evaluation of sensitivity experiments


Table: Sites used in the wind speed verification

Site	Type	Heights (m AGL)
FINO1	Offshore	100, 90, 80, 70, 60, 50, 40, 33 m
FINO2	Offshore	102, 92, 82, 72, 62, 52, 42 m
FINO3	Offshore	106, 100, 90, 80, 70, 60, 50, 40, 30 m
Høvsøre	Coastal	116.5, 100, 80, 60, 40, 10 m
Risø	Land	125, 118, 94, 77, 44 m
Østerild	Land	244, 210, 178, 140, 106, 70, 40, 10 m
Cabauw	Land	200, 140, 80, 40, 20, 10 m
IJmuiden	Offshore	315, 290, 265, 240, 215, 190, 165, 140, 115, 89, 58, 27 m




PBL	Method
MYNN	36 hours simulation, 12 hours spin-up/
YSU	8 days simulation, 1 day spin-up, nudging D1

MYNN, daily MYNL61S1	MYNN, weekly MYNL61W1
YSU, daily MYNL61S1	YSU, weekly YSUL61W1



Phase1: NW WRF sensitivity experiments

MYNN, daily MYNL61S1	MYNN, weekly MYNL61W1
YSU, daily MYNL61S1	YSU, weekly YSUL61W1




Summary statistics
For 8 tall masts in
the North Sea
wind speed at
~100m

Table: NW/MYNL61W1 sir


RUN name	version	note
MYNL61W1	3.6.1	cont
MYNL61W1_V381	3.8.1	sam
MYNL61W1_V391	3.9.1	sam
MYNL61W1_2WAY	3.8.1	two-
MYNL61W1_ERA_0p25	3.8.1	ERA
MYNL61W1_ERA_1p00	3.8.1	ERA
MYNL61W1_MERRA2	3.8.1	MEI

Main differences come
from WRF V3.8.1 vs
V3.6.1




New simulations

Fixes to WRF V3.8.1,
 bl_mynn_mixlength=0,
 COARE OPT=3.0 (sfc
 layer)




Validation of regional simulations: Temporal Variability



Phase2: Optimal domain size


Phase2: NEWA-Light experiments

What is the optimal size of the domains?



Difference from SM long simulations

Annual mean wind speed (m/s), Height: 100 m



Annual mean wind speed bias, $(\bar{U}_{WRF} - \bar{U}_{OBS})/\bar{U}_{OBS}$

site	height (m)	worst sim	bias (%)	best sim	bias (%)
FINO1	100	MD-D1	8.8	SM-L1	7.1
FINO2	102	LG-L1	13.5	SM-L1	9.3
FINO3	100	LG-L1	6.4	SM-L1	2.7
Høvsøre	100	LG-D1	3.2	SM-L1	0.7
Risø	94	LG-L1	10.1	SM-L1	6.8
Østerild	106	LG-L1	16.4	SM-L1	13.0

Smaller domains give better results


but

Is the location of the inflow
conditions that is important?

Phase3: Beta production run

With this in mind...


- Common effort to start simulations waiting for the PRACE application response
- Ten domain NEWA light configuration
 - All domains share the same outer domain, so that the inner grids are coincidental
 - Each European country is contained solely within one domain, except for Sweden, Norway and Finland.
- 13 months (June 2016-June 2017) simulation for BA, FR, GR, IB, IT, SA, and TR at DTU
- 8 years for CE, 8 years for GB at Oldenburg
- 3 years for SB at DTU
- 1 year for IB at CIEMAT



Phase4: Planning for production run

Table 2: Table showing the mesoscale fields and quantities to be served by NEWA.

- 56,700,000 CPU hours granted for the next year at BSC MareNostrum4



Field	Quantity	Time interval means	Heights [m]
Power density	(long-term mean)	Static (all sector sum)	(10,50,75,100,150,200,250, 500
	Sectorwise frequency distribution	Static	10,50,75,100,150,200,250, 500
Horizontal winds	Long-term mean wind speed	Static	10,50,75,100,150,200,250, 500
	Sectorwise wind speed frequency distribution	Static	10,50,75,100,150,200,250, 500
	Sector frequency distribution (wind rose)	Static	10,50,75,100,150,200,250, 500
	Wind speed time series	30-min & 1-monthly	10,50,75,100,150,200,250, 500
	Wind direction time series	30-min & 1-monthly	10,50,75,100,150,200,250, 500
Roughness length	Long-term mean value		Surface
Surface elevation	Long-term mean value		Surface
Friction velocity	Time series	30-min & 1-monthly	
Air temperature	Long-term mean	Static	2,50,75,100,150,200,250, 500
	Time series	30-min & 1-monthly	2,50,75,100,150,200,250, 500
Air pressure	Long-term mean	Static	Surface
	Time series	30-min & 1-monthly	Surface
Air density	Long-term mean	Static	Surface
	Time series	30-min & 1-monthly	Surface
sturbulenc e intensity (TKE as proxy)	Long-term mean	Static	50,75,100,150,200,250, 500
	Time series	30-min & 1-monthly	50,75,100,150,200,250, 500
Specific humidity	Long-term mean	Static	
	Time series	30-min & 1-monthly	Surface layer
Inverse M-O length	Long-term mean	Static	
	Time series	30-min & 1-monthly	Surface layer
Boundary layer height	Long-term mean	Static	-
	Time series	30-min & 1-monthly	-